

UNITED STATES PLANT PATENT APPLICATION

of

L. PERNILLE AND MOGENS N. OLESEN

for

ROSE PLANT NAMED

'POULac013'

## SUMMARY OF THE INVENTION

### BOTANICAL CLASSIFICATION

*Rosa hybrida*

### VARIETY DENOMINATION

5 'POULac013'

The present invention constitutes a new and distinct variety of garden rose plant which originated from a controlled crossing between the female seed parent, an un-  
10 named seedling, and the male parent, an un-named seedling. The two parents were crossed during the summer of 1994 and the resulting seeds were planted in a controlled environment in Fredensborg, Denmark. The new variety is named 'POULac013'.

15 The new variety may be distinguished from its female seed parent by the following characteristics:

The seed parent has a tall growth habit, and has salmon pink flowers. 'Poulac013' is compact and has light pink and apricot blend flower color.

20 The new variety may be distinguished from its male pollen parent by the following combination of characteristics:

The pollen parent has pink flowers while 'Poulac013' has light pink and apricot blend  
25 flower color. Additionally, the pollen parent

has more flower petals than 'Poulac013'.

The objective of the hybridization of this rose variety was to create a new and distinct variety for garden use with unique qualities, such as:

- 5           1.    Uniform and abundant light pink and apricot  
              colored flowers;
2.    Vigorous, but compact and even growth when  
              propagated both as a budded rose and on its own  
              roots;
- 10          3.    Disease resistance.

This combination of qualities is not present in previously available commercial cultivars of this type, known to the inventors, and distinguish 'POULac013' from all other varieties of which we are aware.

15           As part of their rose development program, L. Pernille Olesen and Mogens N. Olesen germinated the seeds from the aforementioned hybridization during winter of 1994-1995 and conducted evaluations on the resulting seedlings in a controlled environment in Fredensborg, Denmark.

20           'POULac013' was selected in the spring 1995 by the inventors as a single plant from the progeny of the aforementioned hybridization.

Asexual reproduction of 'POULac013' by traditional budding and rooted cuttings was first done by L. Pernille  
25           and Mogens N. Olesen in their nursery in Fredensborg,

Denmark in July, 1995. This initial and other subsequent  
asexual propagations conducted in controlled environments  
have demonstrated that the characteristics of 'POULac013'  
are true to type and are transmitted from one generation to  
the next.

#### BRIEF DESCRIPTION OF THE DRAWING

The accompanying color illustration shows as true as  
is reasonably possible to obtain in color photographs of  
this type, the typical characteristics of the buds, flowers,  
leaves, and stems, of 'POULac013'. Specifically illustrated  
in the first drawing;

Fig 1.1; Open flower, stem showing cluster of  
open flowers, branching, and the attachment of  
leaves, buds, and peduncles;

Fig 1.2; Sepals, receptacle, and peduncle;

Fig 1.3; Flower buds at various stages of  
development;

Fig 1.4; Flower petals, detached;

Specifically illustrated in the second drawing;

Fig 2.1; Mature leaves;

Fig 2.2; Bare stems, exhibiting thorns.

### DETAILED DESCRIPTION OF THE VARIETY

The following is a description of 'POULac013', as observed in its growth in a field nursery in Jackson County, Oregon. Observed plants were budded on to *Rosa multiflora* rootstock, and are 3 years of age. Color references are made using the Royal Horticultural Society (London, England) Colour Chart, 1995, except where common terms of color are used.

For a comparison, several physical characteristics of the rose variety 'POULrek', a rose variety from the same inventors described and illustrated in U.S. Plant Patent No. 9,688 issued 12 November 1996, are compared to 'POULac013' in Chart 1.

CHART 1

	'POULac013'	'POULrek'
Petalage	20 petals	25 to 30 petals
Flower diameter	45 to 60	88 to 95 mm
Color of the upper surface of outer petals after flowers open.	White 155D. Yellow Group 6D to 6C.	Red 36D. Yellow 4D petal spot.

## FLOWER AND FLOWER BUD

Blooming habit: Continuous

### Flower bud:

- 5                      Size:                      Upon opening, 18 mm in length  
from base of receptacle to end of  
bud. Average bud diameter is  
10mm.
- 10                      Bud form:                      Pointed ovoid slightly broad  
based.
- Bud color:                      As sepals unfold, petals are  
Red Group 45C to 45D with  
intonations of Red-Purple group  
65A to 63A.
- 15                      Sepals:
- Upper Surface:
- Color:                      Yellow-Green Group 144B.
- Surface:                      Moderately pubescent.
- Lower Surface:
- 20                      Color:                      Yellow-Green Group 144A.  
Marginal anthocyanic  
pigments the color of  
Greyed-Red Group 182A  
observed.
- 25                      Texture:                      Smooth with scant stipitate

glands.

Sepal Shape:

Sepal apex is cirrhose.

Base is flat at union with  
 5 receptacle.

Sepal Margin:

Margins have weak  
 foliaceous appendages on  
 three of the five sepals.

10 Size: 20 mm (l) x 5 mm (w).

Receptacle:

Surface Texture:

Glaucous.

Shape: Funnel shaped.

15 Size: 4 m (h) x 6 mm (w).

Color: Yellow-Green Group 144B.

Anthocyanic pigments the  
 color of Greyed-Red Group  
 181A observed.

20 Peduncle:

Surface: Slightly pubescent with  
 stipitate glands.

Length: 25 to 30 mm average length.

Color: Yellow-Green Group 144D.

25 Anthocyanic pigments the

		color of Greyed-Red Group
		181B observed.
	Strength:	Somewhat strong.
5	Borne:	In clusters of 7 flower buds per stem.
	<b>Flower bloom:</b>	
	Fragrance:	Moderate rose scent.
10	Duration:	The blooms have a duration on the plant on average 10 days. At full maturity, flower petals fall cleanly away from plant.
15	Size:	Flower diameter is 45 to 60 mm when open. Flower depth is 25 to 30 mm in depth.
	Form:	General shape is a deep cup.
	Shape of flower when viewed from the side:	
20	Upon opening, upper part: Flat.	
	Upon opening, lower part: Flat.	
	Open flower, upper part: Flat.	
	Open flower, lower part: Concave.	
25	Petalage:	20 petals on average with 4 petaloids.



Color:

Upon opening, petals:

Outermost petals:

Outer side: Yellow-Orange Group 14D with  
marginal intonations of  
Orange-Red Group 32D and Red  
Group 36D.

Inner Side: White Group 155A with  
intonations of Red Group  
36D.

Innermost petals:

Outer side: Yellow-Orange Group 18D with  
intonations of Orange Group  
32D and Red-Purple Group  
63D.

Inner Side: Yellow-Orange Group 18D with  
intonations of Yellow Group  
8C at middle and basal  
zones.

20            Upon opening, basal petal spots:

Outermost petals:

Outer side: NONE.

Inner Side: Yellow Group 6D to 6C.

Innermost petals:

25                      Outer side:        Yellow Group 6D to 6C.

Inner Side: Yellow Group 6D to 6C.  
 After opening, petals:  
 Outermost petals:  
 Outer side: White Group 155D with light  
 5 intonations Yellow Group 4D  
 at the base and Red Group  
 49D at the margins.  
 Inner Side: White Group 155D.  
 Innermost petals:  
 10 Outer side: Yellow-Orange Group 18D with  
 marginal intonations of  
 Orange-Red Group 32D and Red  
 Group 36D.  
 Inner Side: Yellow-Orange Group 18D.  
 15 After opening, basal petal spots:  
 Outermost petals:  
 Outer Side: NONE.  
 Inner Side: Yellow Group 6D to 6C.  
 Innermost petals:  
 20 Outer Side: NONE.  
 Inner Side: Yellow Group 6D to 6C.  
 General Tonality: Open flowers are Yellow-Orange  
 Group 21B with intonations of  
 25 Orange Group 29B. No change in

5 the general tonality at the end  
of the 10th day. Afterwards,  
general tonality changes to White  
Group 155D with intonations of  
Yellow Group 6D and Red Group  
36D.

**Petals:**

Petal Reflex: Somewhat reflexed.  
Margin: Entire and uniform. Weak  
10 undulations of margin observed.  
Shape: Apex: round.  
Base: acute to round.  
Size: 34 mm (l) x 30 mm (w).  
Texture: Smooth.  
15 Thickness: Thin.  
Arrangement: Not Formal.

**Petaloids:**

Quantity: 3 to 6.  
Color:  
20 Upper Surface: Yellow Group 12D with  
marginal Red Group 36B.  
Lower Surface: Yellow Group 12D with  
marginal intonations of Red  
Group 36B.  
25 Size: 25 mm (l) x 20 mm (w).

**Reproductive Organs:**

**Pistils:**

5                      Length:     5 mm.  
                         Quantity: 42 (actual count).

**Pollen:**

                         Color:     Yellow-Orange Group 17A.  
                         Quantity: Average.

**Anthers:**

10                      Size:        2 mm in length.  
                         Color:     Yellow-Orange Group 17A.  
                         Quantity: 103 (actual count).

**Filaments:**

15                      Color:     Yellow Group 13A.  
                         Length:    8 mm.

**Stigmas:**                      Inferior relative to the  
   filament length and height  
   of the anthers.

                         Color:     Yellow Group 9C.

20                      **Styles:**

                         Color:     Yellow-Green Group 154C.

                         Other Intonations:

   Greyed-Red Group 18D.

25                      **Hips:**        None Observed in the field nursery in  
                         Jackson County Oregon.

## PLANT

5                   **Plant growth:** Compact, upright to bushy. When grown  
as a budded field grown plant on *Rosa multiflora* understock, the average  
height of the plant is 81 cm and the  
average width is 65 cm.

### Stems:

10                   **Color:**  
                    Young wood:       Yellow-Green Group 144B.  
                    Older wood:       Yellow-Green Group 144B.  
                    **Surface Texture:**  
                    Young wood:       Smooth.  
15                   Older wood:       Smooth.

### Thorns:

**Incidence:**       5 thorns per 10 cm of stem.  
                    **Size:**           Average length: 10 mm.  
                    **Color:**          Greyed-Yellow Group 161A.  
20                   **Shape:**          Concave.

**Plant foliage:**   Normal number of leaflets on  
normal leaves in middle of the  
stem: 5 to 7 leaflets.  
                    **Compound leaf size:**       110 mm in length  
25   by 70 mm wide.

5

10

15

20

25

Group 148C.

Stipules:

Quantity: 2 per compound leaf.

Color: Yellow-Green Group 144B.

Length: 25 mm.

Color: Yellow-Green Group 144B.

Rachis:

Length: 50 mm.

Above:

Color: Yellow-Green Group 144B.

Underneath: Thorns.

Leaflet:

5                      Size:                      35mm in length by 30mm wide.

Edge:                      Serrated.

General Shape: Ovate.

Base Shape:              Acute.

Apex Shape:              Round.

10                      Thickness:              Thick.

Glossiness:              Moderate.

Arrangement:            Odd pinnate.

Venation:                Reticulate.

15                      **Disease resistance:**

Above average resistance to mildew, rust, black spot, and Botrytis under normal growing conditions in Jackson County, Oregon.

20                      **Cold Hardiness:**

The variety 'POULac013' has been found to be cold tolerant to USDA Cold Hardiness Zone 6.